C
D

HEBHK
10
Tahun

TB infection among household contacts
Contact tracing is the most effective way to detect TB infection
Before starting TB preventive Treatment (TPT), TB disease must be excluded!
Giving TPT to someone who has TB disease can delay resolution of disease and favour the emergence of drug resistance.
To exclude TB disease, a child should be screened for the following:
i. Symptoms** suggestive of TB disease, and
ii. Chest X-ray
iii. Tuberculin skin test (TST) or IGRA (test needed for children ≥ 5 year old)
Sedekah HEBHK
Warisan Sedekah Melestari Kesejahteraan Masyarakat

World Health Organization
D TB
ANTI TB DRUGS
PHARMACY DEPARTMENT
Selamat Datang ke
PROGRAM KENALI UBAT ANDA
TANYA AHLI FARMASI ANDA
PROGRAM PERKHIDMATAN FARMASI KEMENTERIAN KESIHATAN MALAYSIA
1800 88 6722
JADUAL PENGGUNA UBAT YANG BETUL
WAKTU PENGAMBILAN UBAT YANG BETUL
Dos Sekali Sehari - Setiap 24 jam
Dos 2 Kali Sehari - Setiap 12 jam
Dos 3 Kali Sehari - Setiap 8 jam
Dos 4 Kali Sehari - Setiap 6 jam
TANYA AHLI FARMASI ANDA
1800 88 6722
KENALI UBAT ANDA
Dimanakah anda melupuskan ubat?
Jangan lupuskan ubat di sini
Ubat dalam simpanan pesakit boleh dilupuskan melalui inisiatif
MyMediSAFE
5 Langkah Mudah Pelupusan Ubat
TANYA AHLI FARMASI ANDA
PROGRAM PERKHIDMATAN FARMASI KEMENTERIAN KESIHATAN MALAYSIA
1800 88 6722